



Tikrit University  
College of Veterinary Medicine

# Avian Pox

Subject name: Poultry Diseases

Subject year:2024

Lecturer name: Nawar Ali Jasim

Academic Email:pdvet10@tu.edu.iq



SCAN ME

Lecturers link

## **6-Avian Pox (Fowl Pox) (F.P)**

\***Definition:** - It is highly **contagious** disease of birds characterized by **cutaneous eruption** on **unfeathers** parts of body and or **diphtheritic membrane** of mucous membrane of **upper digestion and Respiratory tract**.

\***Etiology:** - It is caused by avian **pox** virus, belong to pox viridae family, **6** strains recognized:-

**1-Fowl Pox Virus (FPV).**

**2-Turkey Pox Virus (TPV).**

**3-Pigeon Pox Virus (PPV).**

**4-Canary Pox Virus (CPV).**

**5-Sparrow Pox Virus (SPV).**

**6-Quail Pox Virus (QPV).**

These viruses has **tropism** of **epithelial of skin** and **mucous membrane** producing **oval cytoplasmic inclusion body (Bollinger bodies)**, it propagated in CAM and produced characterized grayish foci.

\***Susceptibility:**-

1-Chicken, turkey, pigeon, canary, Quail, pheasant, sparrow and some other wild bird are susceptible, **all ages** and sex are susceptible.

2-Duck and geese are almost **resistance** to natural infection.

\***Mode of Infection:**-

1-Infection occurs through **injured or lacerated skin** and mucous membrane in **head**.

2-Infection can occur by direct contact with **infected bird**.

3-Infection can occur by **bite** of mosquito, mite and other insects.

4-Virus show **resistance** to environmental condition.

\***Incubation Period:** Vary from (4-10) days.

### **\*Clinical Signs:-**

The disease may occur in **4 Forms:-**

- 1-Dry Form (skin form, cutaneous form).**
- 2- Moist Form (wet form, diphtheritic form).**
- 3-Occular Form.**
- 4-Mixed Form.**

#### **1-Dry Form (skin form, cutaneous form).**

Characterized by the appearance **wart like nodular** lesions on the **comb, wattle, eyelids**, and other **nonfeathered** areas of the body.

#### **2- Moist Form (wet form, diphtheritic form).**

1-Characterized by diphtheric **yellowish lesions** occur on the **mucous membranes** of the **mouth, tongue, esophagus, or trachea** with mild or severe **respiratory signs** and interfere with **eating, drinking, and breathing**.

2-As well as there are drop in **egg production** in layer.

**3-Occular Form.** The lesion may found in **eyelids** and cause **complete closure** of both eyes, lead to **starvation and death**.

### **\*Morbidity and mortality:-**

If there are severe infection, the morbidity **10-90%**, and mortality **10-40%**.

### **\*Gross lesion:-**

#### **1-Dry Form (skin form, cutaneous form).**

1-The **characteristic lesion** of the cutaneous form of pox in chickens is a **local epithelial hyperplasia** involving epidermis with formation of **nodules** that first appear as **small white foci** and then rapidly **increase** in size and become **yellow**.

2-**Papules** are formed by the 5th-6th day, this is followed by **vesicular stage**, with formation of extensive **thick lesions**, and the lesions may coalesce and become **rough and gray or dark brown**.

3-After **2 weeks** or sometimes sooner, lesions have areas of inflammation and formation of a **scab**.

#### **2- Moist Form (wet form, diphtheritic form):-**

- 1-In the diphtheritic form, slightly elevated, **white opaque nodules or yellowish patches** develop on the **mucous membranes** of the mouth, esophagus, tongue, or upper trachea.
- 2-**Nodules** rapidly increase in size and coalesce to become a **yellow necrotic pseudodiphtheritic or diphtheritic membrane**, if the membranes are removed, they **leave bleeding erosions**.
- 3-The inflammatory process may extend into **sinuses, pharynx and larynx** and esophagus.
- 4-Often involvement of the **eyes and eyelids** may accompany the formation of lesions in other areas of skin as well as diphtheritic lesions.

**\* Microscopic lesions:-**

- 1-The most important lesion of infection (whether the lesion is cutaneous, diphtheritic, or from infected CAM) is **hyperplasia** of the **epithelium** and enlargement of cells with associated inflammatory changes.
- 2-Characteristic **eosinophilic A type cytoplasmic inclusion bodies** (Bollinger bodies).
- 3-Histopathologic changes of **tracheal mucosa** include initial **hypertrophy and hyperplasia** of **mucus-producing cells** that contain eosinophilic cytoplasmic inclusion bodies.

**\* Diagnosis:-**

- 1-Clinical signs.
- 2- Blood examination (Wright's stain).
- 3-Neutrilization, ELISA.
- 4-Virus isolation.

**\* Differential Diagnosis:-**

- 1-Vitamine A.
- 2-Trichomonas gallinae.
- 3-ILT.
- 4-Pantothenic acid or Biotin deficiency.

**\* Control:-**

First vaccination at (4th-8th) weeks.  
Revaccination at (16-18) weeks.

Referens:

1-Saif, Y. M. (2009). *Diseases of poultry*. Twelfth edition. Iowa. Blackwell.2009. 291-309.