



Tikrit University  
College of Veterinary Medicine

# Lect.6: Microbiology

Subject name: *Campylobacter*

Subject year: Third-year

Lecturer name:

Assist.Lecturer. Hanen omar

Academic Email:



SCAN ME

Lecturers link

Species : *Campylobacter*

Campylobacter fetus subsp. venerealis

Campylobacter fetus subsp. fetus

Campylobacter jejuni subsp. jejuni

Morphology and staining :-

- 1- The shape of these bacteria is like a comma or s-shape or seagull shaped.
- 2- Arrangement : single, pairs, or short chains
- 3- Gram-negative
- 4- Motile/ linear movement or corks crew motility by monotrichous flagella
- 5- Non-spore forming
- 6- Non-capsulated

Cultural characteristics :-

- 1- microaerophilic bacteria
- 2- grows well in the presence of 5% O<sub>2</sub>, 10% CO<sub>2</sub> and 85% N<sub>2</sub>
- 3- requires serum or blood in the primary isolation
- 4- colonies on blood are circle, high, opaque, glistening with regular ridges
- 5- grows on MacConkey agar

Biochemical test :-

Species	Catalase	Nitrate reduction	Lead acetate H <sub>2</sub> S production	Triple sugar iron H <sub>2</sub> S production	Nalidixic acid	cephalothin
<u>C.fetus subsp. venerealis</u>	+	-	-	-	resistant	S
<u>C. fetus subsp. fetus</u>	+	-	+	-	resistant	S

<u>C. jejuni</u> <u>subsp.</u> <u>jejuni</u>	+	-	+	-	susceptible	R
--	---	---	---	---	-------------	---

Diagnosis :-

- 1- isolation of these bacteria from aborted fetus, infected cows and bulls.
- 2- Smear are stained with Gram stain to identify the shape of bacteria
- 3- Antibody florescent test.
- 4- Specimen from fetal absomal contents or vaginal discharge or seminal fluid are cultured on blood agar in atmosphere containing 5% O<sub>2</sub>, 10% Co<sub>2</sub> and 85% N<sub>2</sub>.

## **Brucella**

Species :-

Brucella abortus

Brucella melitensis

Brucella suis

Brucella canis

Brucella ovis

Brucella neotomae

Morphology and staining :-

- 1- Gram -ve
- 2- rod or coccobacilli
- 3-arrangement : single or pairs or small groups or short chains.
- 4-resistant for decolorlization with 0.5% acetic acid while staining with modified Ziel-Nelsen stain or by Koster stain( the bacteria appear red on a blue ground
- 5- Non-motile
- 6- Non-spore forming

Cultural characteristics :-

- 1- Aerobic bacteria except Brucella abortus which requires 5-10% Co<sub>2</sub> for growth.
- 2- On nutrient agar : the colonies of Brucella abortus, Brucella melitensis and Brucella neotomae appear as pin point, circular and translucent.
- 3- On blood agar : the colonies of Brucella abortus, Brucella melitensis and Brucella neotomae appear as pin point, circular, glistering. Non-hemolytic and pale yellow.
- 4- Selective media for Brucella
- 5- Serum agar
- 6- Albimi medium
- 7- Liver extract serum agar
- 8- Serum dextrose agar

note

mixture of antibiotics should be added to the specific media of Brucella, to prevent the growth of fungus and bacterial contamination. The antibiotics include: sulphate, polymyxin B, bacitracin, cycloheximide, Nalidixic acid, Nystatin and Vancomycin.

Diagnosis :-

- 1) rapid presumptive diagnosis by making smear from placental tissue or fetal stomach contents or uterine discharges and staining with modified Zielh-Nelsen stain or Koster stain.
- 2) Bacterial isolation: culturing the samples on the selective media and incubate for 2-3 days, after that tests the colonies by agglutination with antisera.
- 3) Guinea pigs injection
- 4) Serological test including :
  - a- agglutination test
  - b- rapid test ( plate test)
  - c- card test ( rose Bengal plate test)
  - d- milk ring test
  - e- complement fixation test
  - f- florescent antibody techniques

milk ring test:- Brucella antigen staining with hematoxylin, add one drop of Brucella antigen to each of 1 ml of the milk and incubate at 37C° for 30-60 minutes the result would be positive with the formation of blue ring on the top of the test tube.

